INDEX TO VOLUME XI

New species in blackfaced type.

Abies, balsamea, Polyporus picipes on wood of, 323; concolor, Fomes applanatus on wood of, 324; concolor, Fomes pinicola on, 326; concolor, Polystictus versicolor on wood of, 321; concolor, Schizophyllum commune on wood of, 320; concolor, Stereum hirsutum on wood of, 320; grandis, host of Polyporus dryadeus 99, 323; grandis, Thelephora terrestris on seedlings of, 142

Abstracts of papers presented at the, twelfth annual meeting of the American Phytopathological Society, Chicago, December 28 to 31, 1920, 31–60; fourth annual meeting of the Pacific Division, Seattle, June 17–18, 1920, 103–106; second annual meeting Canadian Division, Guelph, Ontario, December 9 to 10, 1920, 202–204

Acacia, sp., host of Fomes applanatus, 326; sp., Schizophyllum commune on, 325; melanoxylon, Schizophyllum commune on, 325.

Acer, sp., host of Cytospora chrysosperma, 157; macrophyllum, Thelephora terrestris on, 142; negundo, Thelephora terrestris on, 142; platanoides, Schizophyllum commune on, 325; rubrum Schizophyllum commune on, 325;, saccharum, Schizophyllum commune on 325; saccharinum, host of Polyporus gilvus, 325

Acid phosphate, effect on control of tomato leaf-spot, 433–445

Acids, tolerance of bacteria to, 244–250

Actinomyces scabies, 6–9, 13–17

Adams, J. F. Observations on wheat scab in Pennsylvania and its pathological histology, 115–124

Adenostegia sp., host of Peridermium filamentosum, 45; host of Peridermium harknessii, 45; host of Peridermium stalactiforme, 45

Aecia, of Puccinia albiperidia borne internally, 343–344

Aesculus californica, Polystictus hirsutus on wood of, 321

Agropyron repens, host of Claviceps sp., 42

Airplane timber, decay of, 45

Air temperature, relation to plant diseases, 446–458

Alfalfa, dry rot of roots of, by Fusarium, 383

Allium cepa var. bulbellifera, host of Botrytis sp., 87–90; host of Puccinia asparagi 87–90

Almond, host of Cyphella marginata, 105

Alnus incana, host of Lenzites saepiaria, 324

Alternaria, sp., on wheat grains, 37; brassicae nigrescens on cantaloupe, 137; **nelumbii** n. sp., on Nelumbium speciosum, 135–140; solani, on tomatoes, 379–380

Amaranthus retroflexus, host of cucurbit mosaic, 47

Amelanchier cusickii, host of **Sclerotinia gregaria**, 106, 225–226

American Phytopathological Society, abstracts, twelfth annual meeting of, 31–60; abstracts, second annual meeting of Canadian Division of, 202–204; report twelfth annual meeting of, 194–201; notice of summer meeting of, 376; fourth annual meeting of Pacific Division of, 103–106

Ammonia, use in seed treatment, 470–471

Anaerobic bacteria in plant tissues, 344

Angular leaf-spot, of cucumber, 297–299

Anthracnose, of cucumber, 298–299; of mango, 26; of garden pea, 500–503

Aphids, carriers of sugar beet mosaic, 349–365

Aplanobacter stewarti on corn, 31

Apothecia, of Cenangium abietis, 166–169; of Sclerotinia cinerea, 176–177; factors affecting production of Sclerotinia cinerea, 495–499

Apparatus; constant humidity case, 51; inoculation platforms, 81; celluloid cone damp chambers, 81–82; Petri dish inoculations, 82–84; chisel-forceps, 175; cone and cup cultures, 239–241; to regulate air temperature and humidity, 446–458
Apple, bitter pit of, 366–370; collar rot of, 105; Cytospora canker of, 105; moldy core of, 105; root rot by Thelephora galactina, 141; Valsa canker of, 236–243; European canker of, on Pacific slope, 464–468
Armillaria, sp.. cultures of, 154; mellea, on Thuja plicata, 176; mellea on Lupinus arboreus, 391–392
Arrhenatherum elatius, host of Claviceps sp., 42
Asclepias syriaca, host of cucurbit mosaic, 47
Ascochyta, rubi, 328; ruborum, 328
Aspen, see Populus tremuloides
Aspergillus sp., 13
ATWOOD, W. M., Physiological studies of the effects of formaldehyde on wheat, 103
Bacillus, apivorus on celery, 55; atrosepticus on potato, 104; carotovorus, 55; coli, 248; melanogenus on potato, 104; phytophthorus on potato, 104, solanisaprus on potato, 104, tracheiphilus on cucumber, 299–300; typhosus, 248; solanacearum, relation of air temperature to, 446–458
Bacteria, sp., causing witches'-brooms of Pinus maritima, 27; on lima bean, 52; on tomato, 55, tolerance to acids in media, 244–250; anaerobic, in plant tissues, 344
Bacterial, blight, of bean, 61–69; blight, of pea, 104; rot of corn, 74–79; spot of tomato 55
Bacteriosis, of celery, 55
Bacterium, angulatum, tolerance to acid, 244–250; campestre, tolerance to acids, 244–250; carotovorus, tolerance to acids 244–250; glycineum, on soybean, 18–24; glycineum, tolerance to acids, 244–250; lacrymans, on cucumber, 298; phaseoli, on bean, 61–69; sojae, on soybean, 18–24, 52; sojae, tolerance to acids, 244–250; tabacum, tolerance to acids, 244–250; tabacum, relation of air temperatures to, 455–456
BAILEY, D. L., see FRASER, W. P.
Barberry, eradication of, 4; host of Puccinia graminis, 39–40
Barium water Bordeaux, 184, 192
Barley, resistance to Helminthosporium sativum, 405–411
BEACH, W. S., A Phytophthora crown rot of rhubarb, 55–56
Bean, bacterial blight of, 61–69
Beech, see Fagus sp.
BENNETT, C. W., see YOUNG, H. C.
BETHEL, ELLSWORTH, Notes on the Peridermiums of pines in Colorado and California, 45; and POSEY, GILBERT B. Investigations to determine the identity of a Cronartium on Ribes in California, 46; Report on pinon blister rust in California (title), 105
Bibliographie der Pflanzenschutzliteratur, 377
Biography and portrait of Dr. F. Kölpin Ravn made available, 101–102
BISBY, G. R., Sclerotinia disease of sunflower in Manitoba, 49; The cooperative potato spraying project (committee report), 60; The cooperative potato spraying project: progress report for 1918 1919, and 1920, 178–193
Bitter pit, of apples and pears, 366–370
Black, leafspeck of cabbage, 44; point of wheat, 515; rot of cabbage, 53–54; spot of rose, 128
Blight, bacterial, of bean, 61–69; of corn seedlings, 35–36; bacterial, of pea, 104; of peach blossoms, 290–294; of pear by Botrytis, 105; late, of potato, 94–96
Blister rust, of pinon, 105
BLODGETT, F. M., and FERNOW, KARL Testing seed potatoes for mosaic and leafroll, 58–59
Blossom blight of peach, 43, 290–294
Blue stone, injury to wheat seed, 103
Bordeaux, dust, 184; mixture, 60, 178–193
Botrytis sp., on Allium cepa, 87–90; causing pear blight, 105

Box elder, host of Septoria negundinis, 100
Break-down of tissues, 44
Bremia lactucae, on Lactuca spp., 54
BRIGGS, FRED N., see MACKIE, W. W.
Bromus, inermis, host of Rhyncosporium secalis, 42; mollis, 385
Brown rot, fungus, apothecia of, 176–177; fungus, age of mummies of, 176–177; of peach, 43
BULLER, A. H. R., Upon the ocellus function of the subsporangial swelling of Pilobolus (title), 204; Upon the chemotactic attraction of fungi for slugs (title), 203; The sound made by fungus guns and a simple method for rendering audible the puffing of Discomycetes (title), 203
Bunt, effect of treatment for, on germination of wheat, 469–484
Bur clover, Cercospora leaf spot of, 311–318
Burgundy mixture, 182
BURKHOLDER, WALTER H., The bacterial blight of the bean: a systemic disease, 61-69
BURNS, G. P., Tip-burn and the leafhopper, 56–57
BUTLER, E. J. The Imperial (British) Bureau of Mycology, 100
BYARS, L. P., Notes on the citrus-root nematode, Tylenchulus semipenetrans Cobb, 90-94

Cabbage, black leaf-speck of, 44; black rot of, 53–54; lightning injury of, 97-98; resistance to yellows, 54
CAESAR, L., Practical hints for the young plant pathologist, 203-204
Callistephus chinensis, host of Septoria callistephi, 50
CAMP, A. F., Convenient chambers for inoculation of plants where high humidity and moderate temperatures are required, 510–512
Canada thistle, host of sunflower Sclerotinia, 49
Canker, of apple, 236–243; of citrus, 70–73, 109–114a; of citrus, in Hawaiian Islands, 377; of poplar, 157–165; European, on Pacific Slope, 464–468
Capsicum annuum, host of cucurbit mosaic, 47
Castilleja sp., host of Peridermium filamentosum, 45; host of P. harknessii 45; host of P. stalactiforme, 45
Ceanothus thyrsiflorus, Schizophyllum commune on, 325
Cedrela sinensis, Schizophyllum commune on, 325
Cenangium abietis on Pinus ponderosa, 166–170; piniphilum on Pinus ponderosa and P. contorta, 294–296
Ceratostomella spp. on wood of white pine, scrub pine, western white pine, western yellow pine, shortleaf pine, loblolly pine, red gum, persimmon, aspen, and northern white cedar, 214-224; coerulea, 217; pilifera, 216, 219, 222, 295, pini, 217
Cercospora medicaginis on bur clover, 311–318
Cereal pathologists, summer meeting of, 177
Celery, bacteriosis of, 55; yellows of, 54–55
Celluloid cones as damp chambers, 81–82
Chlamydospores, of Fomes officinalis, 173-175
Chambers, inoculation, 510–512
Chemotaxy, of fungi for snails, 203
Choke cherry, host of Sclerotinia demissa, 106
Chrysomyxa, abietis, 388
CHURCH, MARGARET B., The relation of mosaic disease to pickling of cucumbers, 28–29; The relation of mosaic disease of cucumbers to pickling, 47
Citrus, aurantifolia, host of canker, 70-73, 112; aurantium subsp. intermedia var. natsu daidai, host of canker, 70–73; blast, inoculation chambers for, 510–512; canker, 70–73; canker in Hawaii, 377; chlorosis, relation to soil conditions, 301–305; fruit spots caused by tissue breakdown, 44; limonia, host of canker, 70–73; maxima, host of canker, 70–73; mitis, host of canker,

70–73; nobilis, host of canker, 70–73; nobilis, resistance to canker, 109–114a; nobilis var. deliciosa, host of canker, 109–114a; nobilis var. deliciosa, host of Tylenchulus semipenetrans, 90–93; root nematode, 90–93; sinensis, host of canker, 70–73; trifoliata, host of canker, 70–73

Cladosporium, spp., in cores of apples, 105; fulvum, on tomato, 232

Clavaria stricta, 143

Claviceps, sp., on Arrhenatherum elatius, 42; on Agropyron repens, 42; on Bromus inermis, 42; on Poa pratensis, 42; on rye, 41–42; on wheat, 42; purpurea, on rye, loss from, 41, 285–289

Coleosporium, campanulae, 388; compositarum, 388; ribicola, new hosts for 45; solidaginis, new hosts for, 45

Colletotrichum lagenarium, on cucumber, 298; pisi on Pisum elatius, jomardi and sativum, 500–503

COLLEY, REGINALD H., The effect of incipient decay on the mechanical properties of airplane timber, 45

Collybia velutipes, on Lupinus arboreus, 389–404; on Lupinus chamissonis, 394

Conifer seedlings, methods of inoculating, 80–84

Control, of potato scab, 6–17, 150–151; of root, stalk, and ear rots of corn, 33–35; of wheat bunt, 37–39; of cereal nematodes, 41; of rice straighthead, 42–43; of peach blossom blight, 43; of apple scab, 43–44; of cucurbit mosaic, 47; of sugar beet mosaic, 48; of tobacco wildfire and angular spot, 48–49; of downy mildew of lettuce, 54; of bacterial spot of tomato, 55; of potato wart, 57; of rose mildew, 128; of rose black spot, 128; of stem girdle of spruce transplants caused by excessive heat, 485–490; of tomato leafspot affected by fertilizers and lime, 433–445

Convenient chambers for inoculation of plants where high humidity and moderate temperatures are required, 510–512

COOK, MEL T., Blossom blight of the peach, 43; The blossom blight of the peach, 290–294; Falling foliage, 337–339; Wilting caused by walnut trees, 346; Sunburn and tomato fruit rots, 379–380

COOLEY, J. S., Disinfecting laboratory cutting instruments, 378

COONS, G. H., and NELSON, RAY, Celery yellows, 54–55

Copper, carbonate dust for bunt, 39; dust for potato blight, 180, 185, 193; dust, Sander's for tomato leaf-spot, 229, 233–235; soap dust for tomato leaf-spot, 229–235; soap liquid for tomato leaf-spot, 230–235; sulphate for bunt, 37–38, 469–484; sulphate for Rhizoctonia solani, 59–60; sulphate in soil, effect on control of tomato leaf-spot, 433–445; sulphate in water of soil, effect on Puccinia malvacearum, 460–462

Corn, bacterial rot of, 32–33, 74–79; decayed by Rhizopus nigricans, 279; ear rot of, 33–34; Fusarium wilt of, 33; Philippine mildew of, 32; root rot of, 32–35; rots of, 33; rust of, 31–32; seedling blight, 35–36; smut of, 32; stalk-rot of, 32–34; Stewart's disease of, 31

Cornus masculus, host of Rhizopus nigricans, 279

Corrosive sublimate, for Rhizoctonia, 150–151; for seed treatment, 470

Corticium, sp., cultures of, 154; vagum, on sugar beet, 48; vagum, temperature influencing pathogenicity of, 56

COTTAM, WALTER P., A "dry rot" disease of alfalfa roots caused by a Fusarium, 383

Cotton, Phoma disease of, 48

Crataegus sp., host of Fomes applanatus, 326

CRAWFORD, R. F., Overwintering of mosaic on species of Physalis, 47

Crocus sativus, host of Rhizoctonia crocorum, 413

Cronartium, sp., on Castilleja, 45; sp., on Quercus, 45; sp., on Ribes, 45; sp., overwintiring of, 45; occidentale, on Pinus monophylla, 46; occidentale, on

Ribes, 46; ribicola, 388; ribicola, investigations of, 170–172
Crown-canker of rose 125–134
Cucumber, angular leaf-spot of, 297–299 anthracnose of, 298–299; bacterial blight, overwintering of, 299–300; host of Colletotrichum lagenarium, 298; mosaic of, 28–29, 46–47; mosaic, transmitted to Asclepias syriaca, 47; mosaic, transmitted to Capsicum annuum, 47; mosaic, transmitted to Martynia louisiana, 47; mosaic transmitted to Micrampelis lobata, 47
Cupressus macrocarpa, host of Polyporus cutifractus, 322; Hydnum ochraceum on wood of, 320; Polyporus carbonarius on wood of, 322
Curly leaf of sugar beet, minimum incubation period of, 424–429
Curly top of sugar beet, 350
Currant, host of, Rhizopus nigricans, 279
Cuscuta sp., on onion, 53
Cutting instruments, disinfecting, 378
Cydonia, host of Rhizopus nigricans, 279
Cylindrocladium scoparium, on rose, losses from, 125–134
Cyphella marginata, on almond, 105; on peach, 105;
Cytospora, sp., on apple, 105; chrysosperma, on poplar, 45, 157–165; chrysosperma, on Acer spp., 157; chrysosperma, on Populus tremuloides, 157–165; chrysosperma, on Populus trichocarpa, 157; chrysosperma, on Populus grandidentata, 157–165; chrysosperma, on Prunus demissa, 157; chrysosperma, on Salix lasiandra, 157; chrysosperma, on Sambucus glauca, 157; chrysosperma, on Sorbus scopulina, 157; leucostoma, on apple, 237

Daedalea, cultures of, 154; unicolor on Salix sp., 326
Damping-off, direct inoculation, 80–84
DANA, B. F., Two new species of Sclerotinia, 106; Two new Sclerotinia diseases, 225-228
Datura stramonium, host of tomato mosaic, 47
Decay, of airplane timber, 45; of sweet potatoes by Rhizopus spp., 279-284
Dialonectria depallens, on Lupinus arboreus, 390
Diabrotica vittata, carrier of Bacillus tracheiphilus on cucumber, 299–300
Diaporthe lupini on Lupinus arboreus, 390
DICKSON, B. T. Studies on mosaic, 202
DICKSON, JAMES G., The influence of soil temperature on the development of the seedling blight of cereals caused by Gibberella saubinetii, 35–36
DICKSON, JAMES G., JOHANN, HELEN and WINELAND, GRACE, Second progress report on the Fusarium blight (scab) of wheat, 35
Diospyrus virginiana, wood stained by Ceratostomella spp., 214–224; wood stained by Lasiosphaeria pezizula, 214–224
Diplodia, sp., 33; rapax on Hevea brasiliensis, 46; lupini on Lupinus arboreus, 390
Direct inoculation, methods of, 80–84
Discomycetes, puffing of spores, 203
Disease, Fiji in Philippine, 251–252
Diseases, plant, relation to air temperature, 446–458
Disinfection, 6–17; of cutting instruments, 378
Dodder, on onion, 53
DOOLITTLE, S. P., Influence of temperature on the development of mosoaic-diseases, 46–47; The relation of wild host plants to the overwintering of curcurbit mosaic, 47; Overwintering of the bacterial wilt of cucurbits, 299–300; see JONES, L. R.
Douglas fir, decayed by Fomes officinalis, 173
Downy mildew, of lettuce, 54; of sugar cane, 371–375; of maize, 372
DRECHSLER, CHARLES, Occurrence of Rhyncosporium on Dactylis glomerata and Bromus inermis, 42
Drought, causing falling foliage of trees, 338-339
Dry rot, of alfalfa roots by Fusarium, 383
DUDDLESTON, B. H., and HOFFER, G. N. The improved rag-doll germinator as

an aid in controlling root, stalk, and ear-rots of corn, 33
DUFF, G. H., Development of the Geoglossaceae (title), 203
DUFRENOY, JEAN, Anaerobic bacteria in plant tissues, 344; Witches'-brooms of Pinus maritima, 27

Early blight, of potato, 182–193
Effect of treatment for bunt on the germination of wheat, 469–484
Eggplant, host of Phytophthora infestans, 504–505
ELLIOTT, JOHN A., A new Phoma disease of cotton, 48; A mosaic of sweet and red clovers, 146–148
ENLOWS, ELLA M. A., and RAND, FREDERICK V., A lotus leaf-spot caused by Alternaria **nelumbii**, sp. nov., 135–140
Epitrix cucumeris, 95
Eradication, of barberry, 4; of milkweed, 47; of wild cucumber, 47
Ergot of rye, losses from, 285–289
ERIKSSON, JAKOB, The mycoplasm theory—is it dispensable or not, 385–388; The life of Puccinia malvacearum Mont. within the host plant and on its surface, 459–463
ERWIN, A. T., see GILMAN, J. C.
ETTER, BESSIE E., Field cultures of wood-rotting fungi on agars, 151–154
Eucalyptus globulus, Polyporus schweinitzii on wood of, 323; host of Fomes applanatus, 326
European canker on the Pacific Slope, 464–468
Eutettix tenella, carrier of sugar beet curly leaf, 424–429
EVANS, NEVADA S., Black point of wheat, 515
Exobasidium sp., on Vaccinium parvifolium, 106
EZEKIEL, WALTER N., Some factors affecting the production of apothecia of Sclerotinia cinerea, 495–499

Fagus, sp., host of Thelephora terrestris, 141; atropunicea, wood discolored by Lasiosphaeria pezizula, 214–224; atropunicea, host of Fomes applanatus, 326
Falling foliage, of trees, 337–339
False mildew of tobacco from Dutch East Indies, 430–432
FARIS, JAMES A., Violet root-rot caused by Rhizoctonia crocorum D.C. in the United States, 412–423
FAULL, J. H. Studies in Fomes fomentarius (title), 203
Fermentation tubes, 21
FERNOW, KARL, see BLODGETT, F. M.
Fertilizers, affecting control of tomato leaf-spot, 433–445
FETHEROLF, N. J., see KORSTIAN, C. F.
Ficus, decayed by Rhizopus nigricans, 279
Fig, see Ficus
Fiji cane disease in Philippines, 251–252, 334–337
Fomes, sp., cultures of 154; annosus, on Thuja plicata, 176; applanatus, on wood of Tsuga heterophylla, 324; applanatus on wood of Picea sitchensis 324; applanatus on wood of Pseudotsuga taxifolia, 324; applanatus on wood of Abies concolor, 324; applanatus on Quercus alba, 325–326; applanatus on Quercus rubra, 326; applanatus on Quercus agrifolia, 326; applanatus on Fagus atropunicea, 326; applanatus on Eucalyptus globulus, 326; applanatus on Crataegus sp , 326; applanatus on Acacia sp., 326; applanatus on Schinus molle, 326; applanatus on Umbellularia californica, 326; fomentarius, 203; laricis, 45; meliae on living Ginkgo biloba, 324; officinalis, chlamydospores of, 173–175; pini on Thuja plicata, 176; pinicola, 45; pinicola on peach, 105; pinicola on prune, 105; pinicola on Abies concolor, 326; pinicola on Pinus lambertiana, 326; pinicola on Pseudotsuga taxifolia, 326; roseus on Pseudotsuga taxifolia, 326
Fomitiporia, cultures of, 154
Formaldehyde, effect on seed wheat, 103; 469–484; for oat smut, 149–150, 203; for potato scab 6–17, 59; for tobacco wildfire and angular spot, 48–49; injury to wheat seed, 103
Fragaria sp., decayed by Rhizopus nigricans, 279
FRASER, W. P., and BAILEY, D. L , Biologic

forms of wheat stem rust in western Canada, 202
FROMME, F. D., Incidence of loose-smut in wheat varieties, 507–510
FROMME, F. D., and WINGARD, S. A. Treatment of tobacco seed and suggested program for control of wildfire and angular-spot, 48–49
Fusarium, sp., 33, 34, 55; avenaceum, 35, 118; blight of wheat, 35, 115–124; causing celery yellows, 54–55; culmorum, 118; culmorum var. leteius, 35; dry rot of alfalfa roots caused by, 383; metochrorum, 118; moniliforme, 34, 35, 76; oxysporium, 56; roseum 118; subulatum, 118; willkommi imperfect stage of Nectria galligena, 464; wilt of corn, 33

Ganoderma sp., cultures of, 154
Garden pea, anthracnose of, 500–503
GARDNER, MAX W., and KENDRICK, JAMES B., Bacterial spot of tomato, 55; and GILBERT, W. W., Field tests with cucumber angular leaf-spot and anthracnose, 298–299; see Kendrick, James B.
GARRETT, A. O., Septoria negundinis Ellis & Ev. in Zion National Park, 100
Geoglossaceae, development of, 203
Gibberella saubinetii, 35, 36, 37; on wheat, 115–124; on corn seedlings, 35–36
GIDDINGS, N. J., Symposium on dusting at the Chicago meeting, 155
GILBERT, W. W., Sclerotium rolfsii on velvet beans, 378; see GARDNER, MAX W.
GILMAN, J. C., A Fusarium wilt of corn in Iowa in 1920, 33; and ERWIN, A. T., Greenhouse propagation of cabbage resistant to yellows, 54; see MELHUS, I. E.
GLOYER, W. O., Septoria leaf blight on China aster, 50–51
GODFREY, G. H., and HARVEY, R. B., Motion pictures of zoospore production in Phytophthora. 145–146
Gortyna immanis attacking Lupinus arboreus, 390
GUSSOW, H. T., Leaf curling in tomatoes, 380–383
Gymnosporangium macropus, internal aecia of 344

HARTER, L. L., WEIMER, J. L., and LAURITZEN, J. I., The decay of sweet potatoes (Ipomoea batatas) produced by different species of Rhizopus, 279–284
HARVEY, R. B., see WEISS, FREEMAN; see GODFREY, G. H.
HASKELL, R. J., Phytophthora infestans on eggplant in the United States, 504–505
HAYES, H. K., and STAKMAN, E. C., Resistance of barley to Helminthosporium sativum P. K. B., 405–411
Head smut of corn in Washington, 515
HEALD, F. C., Relation of spore load to the per cent of stinking smut (Tilletia tritici), 103–104; The skin spot (Oospora pustulans) of the Irish potato, 104–105; Some new hosts for the Rhizoctinia disease, 105; Moldy core of the Stayman Winesap, 105; The relation of spore load to the per cent of stinking smut appearing in the crop, 269–278
Heart rot, of peach trees, 105; of prune trees, 105;
Helminthosporium, sp., 2, 3, 37; avenae, 37; gramineum, 37; interseminatum, 37; sativum on barley, 405–411; sativum causing "black point" of wheat, 515
Hepialus sequoiolus, attacking Lupinus arboreus, 390
Herpotrichia pezizula, 214
HESLER, L. R., Review of ERWIN F. SMITH'S An introduction to bacterial diseases of plants, 107–108
Heterodera radicicola, 93
Hevea brasiliensis, lightning injury of, 46
Hints for the young plant pathologist, 203–204
HOFFER, G. N., see DUDDLESTON, B. H., see TROST, JOHN F.; see SMITH, G. M.
Hollyhock rust, its life within the host plant and on its surface, 459–463
HOPKINS, E. F., Growth and germination of Gibberella saubinetii at varying hydrogen-ion concentrations, 36; Hydrogen-ion concentration of the soil

and seedling infection by Gibberella saubinetii, 36–37; Studies on the Cercospora leaf spot of bur clover, 311–318; Note on the hydrogen-ion concentration of potato dextrose agar and a titration curve of this medium with lactic acid, 491–494
Hopper burn of potato, 188–190
Hordeum vulgare, host of Helminthosporium sativum, 405–411
Hot air, seed treatment by, 471
Hotson, J. W., Collar rot on apple trees in the Yakima Valley (title), 105; A new species of Exobasidium, 106
Hottes, Chas. F., A constant humidity case, 51
Hot water treatment, for citrus root nematodes, 93; of seed corn, 470
Howitt, J. E., Experiments with Haskell's method, or the so-called dry formaldehyde treatment for the prevention of oat smut, 203
Hubert, Ernest E., A chisel forceps, 175; Notes on sap stain fungi, 214–224
Humidity regulating apparatus, 51, 446–458
Humphrey, H. B., see Whetzel, H. H.
Hungerford, Chas. W., A modification of the concentration formaldehyde method of seed treatment, 149–150
Hydnum, sp., cultures of, 154; ochraceum, on trunk of Cupressus macrocarpa, 320
Hydrated lime, for spraying potatoes, 182
Hydrogen-ion, 18–24, 36, 244–250; concentration of potato dextrose agar, 491–494
Hymenochaete tabacina, on wood of Tsuga canadensis and Sequoia sempervirens, 320
Hymenula lupini on Lupinus arboreus, 391
Hyoscyamus niger host of Peronospora hyoscyami, 430
Hypochnus sp., in cores of apples, 105

Imperial (British) Bureau of Mycology, 100
Incidence of loose smut in wheat varieties, 507–510
Inoculation, methods with damping-off fungi, 80–84; chambers, 510–512
Internal aecia, of Gymnosporangium macropus, 344; of Nigredo caladii, 344; of Puccinia albiperidia, 343–344; of Puccinia angustata, 344; of Puccinia cari-bistortae, 344; of Puccinia graminis, 344
Ipomoea batatas, decay by Rhizopus spp., 279–284
Irish potato, see potato
Iron, haematoxylin, 27; sulphate for spraying potatoes, 182.
Irpex, sp., cultures of, 154
Iva sp., killed by sunflower Sclerotinia, 49

Jackson, H. S., and Mains, E. B., The aecidium of the orange leaf rust of wheat, Puccinia triticina, 40; see Mains E. B.
Jennison, Harry Milliken, Observations upon the bacterial blight of field and garden peas in Montana, 104; Bacillus atrosepticus van Hall, the cause of the blackleg disease of Irish potatoes, 104
Johann, Helen, see Dickson, James G.
Johnson, A. G., Recognition of scientific accomplishments, 105; and Leukel, R. W., The nematode disease of cereals, 41; see McKinney, H. H.
Johnson, James, Inheritance of disease resistance to Thielavia basicola, 49; The use of sterilized soils in phytopathological research, 51; The reation of air temperature to certain plant diseases, 446–458
Jones, Fred Reuel, and Vaughan, R. E., Anthracnose of the garden pea, 500–503
Jones, L. R., and Williamson, Maude Miller, Bacterial leaf spot of red clover, 50; and Doolittle, S. P., Angular leaf-spot of cucumber, 297–298; and Keitt, G. W., Eugene Washburn Roark, 327; see Walker, J. C.
Juglans, nigra, causing wilting, 346; regia, anaerobic bacteria in diseased roots of, 349
Keitt, G. W., Second pogress report on, apple scab and its control in Wisconsin 43–44; see Jones, L. R.

KENDRICK, JAMES B., and GARDNER, MAX W., Seed transmission of soybean bacterial blight, 340–342; see GARDNER, MAX W.
KIRBY, R. S., see STAKMAN, E. C.
KOCH, ELIZABETH and RUMBOLD, CAROLINE, Phoma on sweet sorghum, 253–268
KORSTIAN, C. F., and FETHEROLF, N. J., Control of stem girdle of spruce transplants caused by excessive heat, 485–490
KULKARNI, G. S., The susceptibility of Dwarf Milo sorghum to smut, 252

Lactic acid, titration of media with, 491–494
Lactuca canadensis, host of Bremia lactucae, 54; ludoviciana, host of Bremia lactucae, 54; sagittifolia, host of Bremia lactucae, 54; scariola var. integrata, host of Bremia lactucae, 54
Larix occidentalis, host of Thelephora caryophyllea, 144; host of Thelephora terrestris, 142
LARUE, CARL D., Lightning injury in Hevea brasiliensis, 46
Lasiosphaeria pezizula causing sap-stain in wood of beech, red gum and persimmon, 214–224
Late blight of potato, results of spraying for, 180–193; relation of air temperature to, 446–458
LAURITZEN, J. I., see HARTER, L. L.
Leaf, curl of tomatoes, 380–383; roll of potato, 58–59; scorch, of strawberry, 44; spot, angular, of cucumber, 297–299; spot, of lima bean, 52; spot, of lotus, 135–140; spot, of tomato, control of, affected by fertilizers, 433–445
LEE, H. ATHERTON, The increase in resistance to citrus canker with the advance in maturity of citrus trees, 70–73; see MCLEAN, FORMAN T.; see LYON, H. L.
LEHMAN, S. G., Soft rot of pepper fruits, 85–87
Lenzites, sp., cultures of, 154; saepiaria on living peach and prune trees, 105; saepiaria on wood of Populus tremuloides, 324; saepiaria on living Alnus incana, 324
LEONIAN, LEON H., Studies on the Valsa apple canker in New Mexico, 236–243
Leptosphaeria circinans, 414
Lettuce, host of Bremia lactucae, 54
LEUKEL, R. W., see JOHNSON, A. G.
Libocedrus decurrens, host of Polyporus amarus, 176; Stereum hirsutum on wood of, 320; Polystictus versicolor on wood of, 321
Lightning injury, to cabbage, 97–98; to Hevea brasiliensis, 46; to potato, 96–97
Lima bean, bacterial leaf spot of, 52
Lime, for potato spraying, 189, 192; for bunt of wheat, 471–484; following treatment with other chemicals, 471–484; affecting control of tomato leaf spot, 433–445
Lime and sulphur mixture, liquid and dust, for brown rot, 43
Lime-sulphur, for apple scab, 43–44; for potato spraying, 185, 189
LIPMAN, C. B., A contribution to our knowledge of soil relationships with citrus chlorosis, 301–305
Liquidambar styraciflua, see red gum
Literature list of phytopathology, 101
Litmus agar, 20, 22
Loblolly pine, see Pinus taeda
Longleaf pine, see Pinus palustris
Loose smut, incidence in wheat varieties, 507–510
Lotus, leaf spot of, caused by Alternaria **nelumbii** n. sp., 135–140
Low temperature, cause of falling foliage of trees, 338
Lupinus, arboreus, host of Collybia velutipes, 389–404; host of Pleurotus ostreatus, 389–404; host of Armillaria mellea, 391–392; attacked by Hepialus, sequoiolus, 390; attacked by Gortyna immanis, 390; Diplodia lupini on, 390; Stictis lupini on, 390; Diaporthe lupini, on, 390; Dialonectria depallens on, 390; Sphaeria lupini on, 390; Stictus decipiens on 390; Metasphaeria lathyri on, 390; Pleospora leguminum on, 390; Phoma lupini on, 390; Sphaeropsis lupini on, 391; Hymenula lupini on,

391; Valsa lupini on, 391; Sordaria lanuginosa on, 391
Lupinus chamissonis, host of Colybia velutipes, 394
Lycopersicon esculentum, host of Rhizopus nigricans, 279
LYMAN, G. R., Report of the twelfth annual meeting of the American Phytopathological Society, 194–201
LYON, H. L., and LEE, H. ATHERTON, citrus canker in the Hawaiian Islands, 377
MCALPINE, D., Bitter pit in apples and pears: the latest results in preventive measures, 366–370
MCCLINTOCK, J. A., The control of peach brown rot and curculio, 43; Overwintering of mosaic of annuals, 47
MCCUBBIN, W. A., The present status of the potato wart in Pennsylvania, 58
MCFARLAND, FRANK T., Infection experiments with Claviceps, 41-42; see SEYMOUR, EDITH K.
MCKINNEY, H. H., The so-called take-all disease of wheat in Illinois and Indiana, 37; and JOHNSON, A. G., Wojnowicia graminis (McAlp.) Sacc. and D. Sacc. on wheat in the United States, 505–506
MCLEAN, FORMAN T., and LEE, H. ATHERTON, The resistance to citrus canker of Citrus nobilis and a suggestion as to the production of resistant varieties in other citrus species. 109–114a
MCMURRAN, STOCKTON MOSBY, biographical sketch and portrait, 25–26
MACKIE, W. W., and BRIGGS, FRED N., Chemical dusts for the control of bunt, 38–39
Macrosiphum solanifolii, 95
Macrosporium, sp., on onion, 53; foot rot, of tomato, 29; solani, on tomato, 29
MAINS, E. B., and JACKSON, H. S., Two strains of Puccinia triticina on wheat in the United States, 40; see JACKSON, H. S.
Maize, Sclerospora spontanea on, 372
Malva silvestris, host of Puccinia malvacearum, 463
Marasmius sp., cultures of, 154
Marssonia, potentillae, on strawberry, 44
MARTIN, WILLIAM H., Inoculated vs. uninoculated sulfur for the control of common scab of potatoes, 58
Martynia louisiana, host of cucurbit mosaic, 47
MASSEY, L. M., Experimental data on losses due to crown-canker of rose, 125–134
MEDALLA, MARIANO G., Fiji disease of sugar cane in the Philippine Islands, 251–252
Medicago, arabica, mosaic of, 147; maculata, host of Cercospora medicaginis, 311–318; sativa, 146; sativa, host of Rhizoctonia crocorum, 412
Melampsora salicina, 388
MELCHERS, L. E., Ecologic and physiologic notes on corn smut, Ustilago zeae, 32; Rhizopus sp. associated with a decay of unripe strawberries in the field, 44
MELHUS, I. E., Bremia on hothouse lettuce 54; Cooperative potato seed treatment experiments, 59–60; and GILMAN, J. C., Measuring certain variable factors in potato seed treatment experiments, 6–17
Melilotus alba, mosaic of, 146–148
Mercuric chloride, for potato seed treatment, 6–17, 59–60
Metasphaeria lathyri, on Lupinus arboreus, 390
Micrampelis lobata, 299; eradication of, 47; host of cucurbit mosaic, 47
Micrococcus, phytophthorus, 104
Microsphaera euonymi, 388
Mildew, of rose, 128; downy of sugar cane, 371–375; false, of tobacco from Dutch East Indies, 430–432
Milkweed, eradication of, 47
Minimum incubation period of curly leaf of sugar beet, 424–429
Mistletoe, mucilage of berries an adhesive, 99
Mollisia earliana, perfect stage of Marssonia potentillae, 44
Mollisiose of strawberry, 44
Monilia, see Sclerotinia
MONTEITH, JOHN, JR., Seed transmission and overwintering of cabbage black rot, 53–54

MORRIS, H. E., and SWINGLE, D. B., An important new disease of the cultivated sunflower, 50
MORSE, W. J., The transference of potato blight by insects, 94–96
Mosaic, 202; of cucumbers, 28–29, 47; diseases influenced by temperature, 46–47; carried by Myzus persicae, 349–365; of Physalis longifolia, 47; of potato, 58–59; of sweet and red clover, 146–148; of Trifolium repens, 146; of Vicia faba, 202; of raspberry, 202; of sugar beets, 349–365; of tobacco, 388; of tobacco, relation to air temperature to 446–458; wild host plants of cucurbit mosaic, 47
Mummies, brown rot, 176–177
Mycoplasm theory, is it dispensable or not, 385–388
Mycosphaerella **rubi** on Rubus spp., 328–331; perfect stage of Septoria rubi, 328–333
Myzus persicae carrier of sugar beet mosaic, 349–365

Nectria, cinnabarina, 467; galligena on apple and pear, 464–468
Negundo, host of Septoria negundinis, 100
NELSON, RAY, Tissue breakdown in fruits and vegetables, 44; see Coons, G. H.
Nelumbium speciosum, host of Alternaria **nelumbii** sp. nov., 135–140
Nematode, disease of cereals, 41
Neofabraea melicorticis, 466–467
NEWTON, MARGARET, Biologic forms of wheat rust in Canada, 202
Nicandra physalodes, host of tomato mosaic, 47
Nicotiana glauca, 430
Nigredo caladii, internal aecia of, 344
Northern white cedar, see Thuja occidentalis
Note on the hydrogen-ion concentration of potato dextrose agar and a titration curve of this medium with lactic acid, 491–494

OBERLY, E. R., List of literature on phytopathology, 101
Onion, Cuscuta sp. on, 53; rust of, 87–90; Puccinia asparagi on, 87–90, smut of, 52–58; Macrosporium rot of, 53; Botrytis sp. on, 87–90; Rhizoctonia on, 105
Ophiobolus graminis, 505
Orange leaf rust, 40
Orthocarpus sp., host of Peridermium filamentosum, harknessi and stalactiforme, 45
ORTON, C. R., Lightning injury to potato and cabbage, 96–98; and WEISS, FREEMAN, The reaction of first generation hybrid potatoes to the wart disease, 306–310; see WEISS, FREEMAN
ORTON, W. A., Compilation of list of new fungi, 29–30
OWENS, C. E., see ZELLER, S. M.

PALM, B. T., The false mildew of tobacco introduced into the United States from the Dutch East Indies? 430–432
Paraformaldehyde for seed treatment, 473
PARKER, CHAS. S., Head smut of corn in Washington, 515
Pea, bacterial blight of, 104
Peach, blossom blight of, 43, 290–294; control of brown rot of, 43; host of Cyphella marginata, 105; host of Lenzites saepiaria, 105; host of Fomes pinicola, 105; host of Sclerotinia cinerea, 43, 176–177; host of Trametes carnea, 105
Pear, bitter pit of, 366–370; blight, caused by Botrytis sp., 105; decayed by Rhizopus nigricans, 279; European canker of, on Pacific Slope, 464–468
Pecan, diseases, 26; rosette, 26; wood rot, 26.
Pedicularis sp., host of Peridermium filamentosum, 45; host of P. harknessii 45; host of P. stalactiforme, 45.
Penicillium sp., 13; in cores of apples, 105
PENNINGTON, L. H., SNELL, W. H., YORK H. H., and SPAULDING, Perley, Investigations of Cronartium ribicola in 1920, 170–172
Pepper, host of cucurbit mosaic, 47; fruit rot by Pythium deBaryanum, 85–87
Peridermium, cerebrum on Quercus sp., 45; cerebrum, new hosts of, 45; fila-

mentosum, on Pinus sp., 45; filamentosum, on Castilleja sp., 45; filamentosum, on Orthocarpus sp., 45; filamentosum, on Pedicularis sp., 45; filamentosum, on Adenostegia sp., 45; harknessii, on Pinus sp., 45; harknessi on Castilleja sp., 45; harknessii, on Orthocarpus sp., 45; harknessii, on Castilleja sp., 45; harknessii, on Orthocarpus sp., 45; harknessii, on Pedicularis, sp., 45; harknessii, on Adenostegia sp., 45; stalactiforme, on Adenostegia sp., 45; stalactiforme, on Castilleja sp., 45; stalactiforme, on Orthocarpus sp., 45; stalactiforme, on Pedicularis, sp., 45;

Peronospora, ficariae, 388; hyoscyami, on tobacco, origin in Dutch East Indies? 430–432; hyoscyami, on Hyoscyamus niger, 430; hyoscyami, on Nicotiana glauca, 430; spinaciae, on spinach, 386–388; trifoliorum, 388

Persimmon wood stained, by Ceratostomella sp., 214–224; by Lasiosphaeria pezizula, 214–224

Personals, 30, 102, 177, 252, 348, 383–384, 516

Petri dish cultures, for direct inoculation of conifer seedlings, 82–84

Pholiota sp., cultures of, 154

Phoma, sp., on cotton, 48; insidiosa, on sorghum, 253–268, 513–514; insidiosa, on Sudan grass, 345; lupini, on Lupinus arboreus, 390

Phoradendron, mucilage of berries, 99

Phragmidium, potentillae, 388; subcorticium, 388

Physalis longifolia, host of tomato mosaic, 47

Phytopathological, research in Russia, 156; Society, American, see American Phytopathological Society; Society of France, 100–101; Society in the Philippines, 516

Phytophthora sp., 145–146; cryptogea, 29; foot rot, 29; infestans, 386–387; infestans, transference of, 94–96; infestans, on Solanum melongena, 504–505; infestans, relation to air temperature, 446–458; rot of rhubarb, 55–56; infestans, on eggplant in the United States, 504–505

Picea, engelmanni, host of Polyporus dryadeus, 99; engelmanni, host of Thelephora terrestris, 142; engelmanni, control of stem girdle of, caused by heat, 485–490; excelsa, anaerobic bacteria from tumors of, 344; mariana, host of Thelephora terrestris, 142; rubens, host of Schizophyllum commune 320; sitchensis, host of Polyporus berkeleyi, 323; sitchensis, Fomes applanatus on wood of, 324; sitchensis, Polyporus caesius on wood of, 321; sitchensis, Polystictus versicolor on wood of, 321

Pickering's solution, 194

Pilobolus sp., 204

Pinon blister rust, 105

Pinus, cembra, 171; contorta, host of Cenangium **piniphilum** 294–296; contorta, host of Thelephora terrestris 142; densiflora, 171; divaricata, host of Thelephora terrestris, 142; echinata, wood stained by Ceratostomella sp., 214–224; flexilis, host of Cronartium ribicola, 171; koraiensis, host of Cronartium ribicola, 171; lambertima, host of Fomes pinicola, 326; maritima, witches' brooms of, 27; monophylla, host of Cronartium occidentale, 46; monticola, host of Thelephora terrestris, 142; monticola, wood stained by Ceratostomella sp., 214–224; mugho, 171; palustris, 221; ponderosa, host of Cenangium piniphilum, 294–296; ponderosa, host of Thelephora terrestris, 142; ponderosa, host of Thelephora fimbriata, 143; ponderosa, host of Cenangium abietis, 166–169; ponderosa, host of Fomes officinalis, 173; ponderosa, wood stained by Ceratostomella sp., 214–224; radiata, Stereum hirsutum on wood of, 320; radiata, Polystictus versicolor on wood of, 320; radiata, Polyporus dichrous on wood of, 323; rigida, host of Thelephora fimbriata, 144; sabiniana, Polystictus hirsutus on wood of, 321; strobus, host of Cenangium abietis, 166; strobus, host of Thele-

phora terrestris, 141; strobus, host of Crenartium ribicola, 171; strobus, wood stained by Ceratostomella sp., 214–224; sylvestris, 171; taeda, wood stained by Ceratostomella sp., 214–224; virginiana, wood stained by Ceratotosmella sp., 214–224
Pisum, elatius, host of Colletotrichum pisi, 500–503; jomardi, host of Colletotrichum pisi, 500–503; sativum, host of Colletotrichum pisi, 500–503
Plant disease survey of England and Wales, 376
Plasmodiophora brassicae, 433
Platanus occidentalis, host of Polyporus gilvus, 325
Pleospora leguminum on Lupinus arboreus 390
Pleurotus, sp., cultures of, 154; ostreatus on Lupinus arboreus, 389–404
Plum, host of Sclerotinia cinerea, 176-177; fruit decayed by Rhizopus nigricans, 279
Poa pratensis, 42
Poke, mosaic of, 47
Polychrome-blue-tannin-orange, 27
Polyporus sp., cultures of, 154; amarus, 176; berkeleyi, on Picea sitchensis, 323; caesius, on wood of Picea sitchensis, 321; cutifractus, on Cupressus macrocarpa, 322; dryadeus, on conifers, 99; dryadeus wood of Abies concolor, 323; dryadeus, on wood of Abies grandis, 323; dichrous, on wood of Pinus radiata 323; perennis, 143; picipes, on wood of Abies balsamea, 323; carbonarius, on wood of Cupressus macrocarpa, 322; gilvus, on Acer saccharinum, 325; gilvus, on Platanus occidentalis, 325; gilvus, on Quercus coccinea, 325; gilvus, on Rhus toxicodendron, 325; leucospongia, on wood of Populus tremuloides, 322; nidulans, on wood of Tsuga canadensis, 323; schweinitzii, 45; schweinitzii, on Thuja plicata, 176; schweinitzii, on stump of Eucalyptus globulus, 323; sulphureus, on Thuja plicata, 176
Polystictus, sp., cultures of, 154; abietinus, on wood of Quercus sp., 321; hirsutus, on wood of Pseudotsuga taxifolia, 321; hirsutus, on wood of Pinus sabiniana, 321; hirsutus, on wood of Aesculus californica, 321; hirsutus, on Prunus sp., 325; versicolor, on wood of Pinus radiata, 320; veriscolor, on wood of Picea sitchensis, 321; veriscolor, on wood of Pseudotsuga taxifolia, 321; versicolor, on wood of Abies concolor, 321; versicolor, on wood of Sequoia sempervirens, 321; versicolor, on wood of Tsuga heterophylla, 321; versicolor, on wood of Libocedrus decurrens, 321
POOLE, R. F., Recent studies on bacteriosis of celery, 55
Poplar canker, caused by Cytospora chrysosperma, 45, 157–165
Populus, caroliniana, host of Cytospora chrysosperma, 45; deltoides, host of Cytospora chrysosperma, 163-165; grandidentata, host of Cytospora chrysosperma, 45, 158–165; tremuloides, host of Cytospora chrysosperma, 45, 158–165; tremuloides, wood stained by Ceratostomella sp., 214-224; tremuloides, Polyporus leucospongia on wood of, 322; tremuloides, Lenzites saepiaria on wood of, 324; trichocarpa, host of Cytospora chrysosperma, 157
Poria, sp., cultures of, 154; weirii, on Thuja plicata, 176
PORTE, W. S., see PRITCHARD, FRED J.
PORTER, R. H., Cooperative seed treatment using hot formaldehyde, 59
POSEY, GILBERT B., see BETHEL, ELLSWORTH
Potassium, sulphide for seed treatment, 471; sulphate, effect for control of tomato leaf spot, 435,
Potato, black leg of, caused by Bacillus atrosepticus, 104; cooperative spraying report, 178–193; dextrose agar, hydrogen-ion concentration of, 491–494; early blight of, 180–193; late blight of, spraying for, 180–193; late blight of, transference of, 94–96; late blight of, relation to air temperature, 446–458 leaf roll of, 58–59; lightning injury of, 96–97; mosaic of, 47; mosaic of, testing for, 58–59; scab, 6–17, 58–59; seed

treatment of, 58–60, 150–151; skin spot of caused by Osopora pustulans, 104; spraying of, 178–133; tip burn of, 56–57; wart of, susceptibility tests for 57; wart of, enzyme activity of, 57; wart of, status in Pennsylvania, 58; wart of, heredity of resistance to, 306–310

POVAH, ALFRED H. W., Valsa poplar canker, 45 An attack of poplar canker following fire injury, 157–165

Pre soaking seed before treatment with chemicals, 473–474

PRITCHARD, FRED J. and PORTE, W. S., Use of copper soap dust as a fungicide, 229–235; Effect of fertilizers and lime on control of tomato leaf spot (Septoria lycopersici), 433–445

Production of apothecia of Sclerotinia cinerea, factors affecting, 495–499

Prune heartrot, caused by Fomes pinicola, 105; by Lenzites saepiaria, 105; by Trametes carnaea, 105

Prunus, sp., decayed by Rhizopus nigricans, 279; host of Polystictus hirsutus, 325; demissa, host of Cytospora chrysosperma. 157; demissa, host of Sclerotinia **demissa**, 227–228

Pseudomonas, sp., on red clover, 50; citri, on Citrus spp., 109–114

Pseudotsuga taxifolia, host of Thelephora terrestris, 142; Polystictus versicolor on wood of, 321; Polystictus hirsutus on wood of, 321; Fomes applanatus on wood of, 324; host of Fomes pinicola 326; host of Fomes roseus, 326

Puccinia albiperidia, internal aecia of, 343–344; angustata, internal aecia of 344; asparagi, on onion, 87–90; caribistortae, internal aecia of, 344; chrysanthemi, 388; glumarum, 4, 205, 385; graminis, effect on F_2 generation of wheat (durum X vulgare), 205–213; graminis, internal aecia of, 344; malvacearum, 459–463; Ribis, 388; sorghi, 31–32; suaveolens, 388; tragopogonis, 388; triticina, 40

PUTTICK, G. F., The reaction of the F_2 generation of a cross between a common and a durum wheat to two biologic forms of Puccinia graminis, 205–213

Pyrus malus, decayed by Rhizopus nigricans, 279

Pythium de Baryanum, on pepper fruits, 85–87

Quercus, sp., host of Cronartium cerebrum 45; alba, host of Thelephora terrestris, 142; alba, host of Fomes applanatus, 325–326; alba, Polystictus abietinus on wood of, 321; agrifolia, host of Fomes applanatus, 326; coccinea, host of Polyporus gilvus, 325; rubra, host of Fomes applanatus, 326

Quince, host of Rhizopus nigricans, 279

RAND, FREDERICK V., see ENLOWS, ELLA M. A.

Raspberry, decayed by Rhizopus nigricans 279; mosaic of, 202

RATHBUN, ANNIE E., Methods of direct inoculation with damping-off fungi, 80–84

RAVN, F. KOLPIN, biographical sketch and portrait, 1–5; reprints of biography and portrait, 101–102

Ravn Scandinavian-American fellowship in plant pathology, 348

Razoumofskya, mucilage of berries of, 99

Red gum, wood stained by Ceratostomella sp., 214–224; wood stained by Lasiosphaeria peziziula, 214–224

Red heart of lettuce, 44,

Red raspberry, decayed by Rhizopus nigricans, 279

REDDICK, D. Foot rot of tomato, 29; New abstract Journal Zentralblatt fur die gesamte Landwirtschaft mit Einschluss der Forst- und Teichwirtschaft, der Tier-Pathologie und Medizin., 29

REDDY, CHAS. H., Experiments with Stewart's disease on dent, flint and sweet corn, 31

REINKING, OTTO A., Fiji disease of sugar cane in the Philippine Islands, 334–337

Resistance, of corn to rots, 33–34; of wheat to take-all, 37; of wheat to nematodes, 41; of tobacco to Thielavia basicola, 49; of China asters to Septoria callistephi, 50; of celery to yellows, 54–

55; to bacterial rots, 55; of potatoes to wart, 57; of citrus to canker, increase in resistance with age of tree, 70–73; of Citrus nobilis to canker, 109-114; of potatoes to late blight, 183-184, 185, 193; of strawberries to mollisiose, 202–203; of Dwarf Milo sorghum to Sphacelotheca sorghi and S. cruenti, 252; of sugar cane to Fiji disease, 336-337; of barley to Helminthosporium sativum, 405–411

Rhizoctonia sp. on bean seedlings, 63; on onions, 105; on potato, 105; on strawberries, 105; corrosive sublimate treatment for, 150–151; crocorum on Solanum tuberosum, 412–423; on alfalfa, 417-423; solani, 6-7, 9-17, 56, 59-60, 412, violacea, synonym of R. crocorum, 413; medicaginis, 414

Rhizopus, sp., 13; sp., causing strawberry rot, 44; arrhizus, 281–284; artocarpi, 281-284; chinensis, 281-284; delemar, 281-284; maydis, 281-284; microsporus, 281-284; nigricans, 279–294; nodosus, 281-284; oryzae, 281-284; reflexus, 281-284; tritici, 281-284

RHOADS, ARTHUR S., Some new or little known hosts for wood-destroying fungi. III., 319–326; The pathology of Lupinus arboreus, with special reference to the decays caused by two wound-parasites—Collybia velutipes and Pleurotus ostreatus, 389-404

Rhubarb, Phytophthora rot of, 55–56, 145–146

Rhus, hirta, Schizophyllum commune on, 325; toxicodendron, host of Polyporus gilvus, 325; vernix, Schizophyllum commune on, 325

Rhyncosporium secalis, on Bromus inermis 42; on Dactylis glomerata, 42

Ribes sp., host of Cronartium occidentale, 46; decayed by Rhizopus nigricans, 279; cynosbati, host of Cronartium ribicola, 170; glandulosum, host of Cronartium ribicola, 170–171; glandulosum, host of Puccinia albiperidia, 343–344; prostratum, host of Thelephora terrestris, 142; rotundifolium, host of Cronartium ribicola, 170

Rice, Sclerotium oryzae on, 42; Sclerotium rolfsii on, 42; seedling blight of, 42; stem rot of, 42; straighthead of, 42–43

RICHARDS, B. L., A dry rot of the sugar-beet caused by Corticium vagum, 48; The pathogenicity of Corticium vagum as affected by soil temperature, 56; Pathogenicity of Rhizoctonia and the fungus as a factor in crop production (title), 106; Soil temperature as a factor in influencing pathogenicity of Rhizoctonia with special reference to the potato (title), 106

ROARK, EUGENE WASHBURN, biographical sketch of, 327; The Septoria leaf spot of Rubus, 328–333

ROBBINS, W. W., Mosaic of sugar beets, 48; Mosaic disease of sugar beets, 349–365

ROBERTS, JOHN W., Stockton Mosby McMurran, biographical sketch and portrait, 25–26; The age of brown-rot mummies and the production of apothecia, 176-177

Root rot, violet, caused by Rhizoctonia crocorum, 412–425

Rose, black spot of, 128; crown canker of, 125–134; mildew, 128

ROSEN, H. R., A bacterial root and stalk rot of field corn, 32–33; Further observations on a bacterial root and stalk rot of field corn, 74–79

Rubus sp., decayed by Rhizopus nigricans, 279; sp., host of Septoria **rubi**, 328-331; sp., host of Mycosphaerella **rubi**, 328–331; allegheniensis, host of Mycosphaerella **rubi**, 329; hispidus, host of Mycosphaerella **rubi**, 329; parviflorus, host of Mycosphaerella **rubi**, 329; strigosus, host of Mycosphaerella **rubi**, 329

RUMBOLD, CAROLINE, and TISDALE, ELIZABETH K., Notes on Phoma insidiosa Tass. found on Sudan grass, 345; Phoma insidiosa on sorghum, 513–514; see KOCH, ELIZABETH;

Russian Mycological and Phytopathological Society, 346-347

Rye ergot, 41; loss from, 285-289

Saccharum spontaneum, resistant to Fiji disease, 337
Salix sp., host of Daedalea unicolor, 326; sp., host of Trametes suaveolens, 326; lasiandra, host of Cytospora chrysosperma, 157
Sambucus glauca, host of Cytospora chrysosperma, 157
Scab, of apple, 43–44; of potato, 6–17, 58; of wheat, 115–124
Schinus molle, host of Fome applanatus, 326
Schizophyllum commune, on Picea rubens, 320; on Abies concolor, 320; on Sequoia sempervirens, 320; on Rhus hirta, 325; on Rhus vernix, 325; on Cedrela sinensis, 325; on Acer platanoides, 325; on Ceanothus thyrsiflorus, 325; on Acacia melanoxylon, 325; on Acacia spp., 325
Sclerospora, sacchari, on sugar cane, 371–375; spontanea, on sugar cane, 372; spontanea on maize, 372
Sclerotinia sp., on sunflowers, 49, 50; cinerea, on peach, 43, 176–177, 290-294; cinerea, on plum, 176–177; cinerea, factors influencing production of apothecia of, 495–499; **demissa**, on choke cherry, 106; **demissa**, on Prunus demissa 227–228; fructigena on apples, 495; gregaria, on Amelanchier cusickii, 106 225–226; laxa, on apricots, 495
Sclerotium, oryzae, 42; rolfsii, 42; rolfsii, on velvet bean, 378
Scrub pine, see Pinus virginiana.
Seed, coat cracking of, leading to injury from seed treatment, 469–484; treatment, injury from, 103; treatment, of corn, 35, 37; treatment of potato, 6-17, 59, 150–151; treatment of tobacco, 48–49; treatment of tomato, 55; treatment of wheat, 38–39; treatment of wheat with various chemicals, 469-484; treatment followed by lime bath, 469–484
Septoria, callistephi, attacking Callistephus chinensis, 50; lycopersici, 379; lycopersici controlled by copper soap dust, 229–235; lycopersici, control of, effected by fertilizers and lime, 433-445; negundinis on box elder, 100; rubi on Rubus spp., 328–331, rubi imperfect form of Mycosphaerella **rubi** 328–331
Sequoia sempervirens, Schizophyllum commune growing on wood of, 320; Hymenochaete tabacina on wood of, 320; Polystictus versicolor, on wood of, 321
SEVERIN, HENRY H. P. Minimum incubation periods of causative agent of curly leaf in beet leafhopper and sugar beet, 424–429
SEYMOUR, EDITH K. and MCFARLAND, FRANK T. Loss from rye ergot, 41; 285–289
SHAPOVALOV, M., Phytopathological research in Russia, 156; Russian Mycological and Phytopathological Society, 346–347
Shortleaf pine, see Pinus echinata.
SHUNK, I. V. and WOLF, F. A., Further studies on bacterial blight of soybean, 18–24; Soybean bacterial blight, 52; see WOLF, FREDERICK A.
Sicyos angulata, 299
Silver fir, host of Thelephora terrestris, 141
SLAGG, C. M., A new seedling disease of tobacco, 49
SMITH, G. M. and HOFFER, G. N., Three methods of controlling the root, stalk and ear rots of corn, 34
Smut, head, of corn in Washington, 515; of onion, 52–53; of sorghum, 252; of wheat, 269–278
SNELL, W. H., Chlamydospores of Fomes officinalis in nature, 173–175; see PENNINGTON, L. H.
Sodium, chloride, 37–38; nitrate, effect on control of tomato leaf spot, 433–445
Soil relationships to citrus chlorosis, 301–305
Solanum, dulcamara, host of tomato mosaic, 47; melongena, host of Phytophthora infestans, 504–505; nigrum, host of tomato mosaic, 47; tuberosum, host of Rhizopus nigricans, 279; tuberosum, host of Rhizoctonia crocorum, 412–422; tuberosum, host of Rhizoctonia solani, 412

Solutions of sugar, 21–23
Some factors affecting the production of apothecia of Sclerotinia cinerea, 495–499
Sorbus scopulina, host of Cytospora chrysosperma, 157
Sorghum susceptible to smuts, 252; sweet, host of Phoma insidiosa, 253, 268, 513-514
Sow thistle, killed by sunflower Sclerotinia, 49
Soybean, bacterial blight of, 18–24, 52; seed transmission of bacterial blight, 340-342; blight, caused by Bacterium glycineum, 18–24, 52, 340–342; blight, caused by Bacterium sojae, 18–24, 52, 340–342
SPAULDING, PERLEY, see PENNINGTON, L H.; Plant disease survey of England and Wales, 376; Bibliographie der Pflanzenschutzliteratur, Das Jahr 1920 377
Sphacelotheca, cruenti, 252; reiliana, on corn in Washington, 515; sorghi, 252
Sphaeria lupini, on Lupinus arboreus, 390
Sphaeropsis lupini, on Lupinus arboreus, 391
Sphaerotheca, mors-uvae, 388; pannosa, 388; pannosa var. rosae, on rose, 128
Spore load of seed, relation to smut in crop, 103–104
Spot blotch disease of barley, 405–411
Spray injury causing falling of foliage of trees, 339
Spraying, of potato, 178-193
Spruce, host of Thelephora terrestris, 141
STAKMAN, E. C., KIRBY, R. S. and THIEL, A. F., The regional occurrence of Puccinia graminis on barberry, 39–40; see HAYES, H. K.
Stayman Winesap, moldy core of, 105
Stem, girdle of coniferous seedlings, control of, 485–490; rust of wheat, 202
Stereum, sp., cultures of, 154; hirsutum, on wood of Pinus radiata, Libocedrus decurrens, Abies concolor and Sequoia sempervirens, 320
STEVENS, F. L., Helminthosporium and wheat foot-rot, 37
Stewart's disease, of corn, 31
Stictis, lupini on Lupinus arboreus, 390; decipiens on Lupinus arboreus, 390
Stinking smut, relation of spore load of seed to smut in crop, 269-278
STONE, R. E., Leaf scorch or mollisiose of strawberry, 44; 202–203
Strawberry, fruit decayed by Rhizopus nigricans, 44, 279; host of Marssonia potentillae, 44, 202; host of Mollisia earliana, 44; leaf scorch, 44, 202; mollisiose of, 44, 202; host of Rhizoctonia, 105
Stripe rust, of wheat, 4
Sudan grass, host of Phoma insidiosa, 345
Sugar beet, curly leaf, carried by Eutettix tenella, 424–429; host of Corticium vagum, 48; mosaic of, 48, 349–365; transmitted by aphids, 349–365; carried by Myzus persicae, 349–365
Sugar cane, downy mildew of, in Philippines, 371–375; Fiji disease of, 251–252 334–337; mosaic of, 47; Sclerospora sacchari on, 371–375; Sclerospora spontanea on, 372
Sulphur, dioxide for seed treatment, 470; for potatoes, 185; ground for rose mildew, 128; inoculated for potato scab, 58
Sulphur and lime, dust for brown rot, 43; spray for brown rot, 43
Sulphuric acid for seed treatment, 470
Summer field meeting of cereal pathologists, 177
Sunburn of tomatoes, 379–380
Sunflower, Sclerotinia disease of, 49–50
Sunscald causing falling foliage of trees, 338–339
Susceptibility of Dwarf Milo sorghum to smuts, 252
Sweet potato, decay by Rhizopus spp., 279–284
Sweet sorghum, host of Phoma insidiosa, 253–268
SWINGLE, D. B., see MORRIS, H. E.
Symposium on dusting, 155
Systemic disease, bacterial bean blight, 61-69

TAYLOR, MINNIE W., Internal aecia of Puccinia albiperidia Arthur, 343-344

Temperature, affecting pathogenicity of Corticium vagum, 56; regulating apparatus, 446–458; air, relation to plant diseases, 446–458
Thalictrum sp., host of Puccinia triticina, 40
Thelephora, caryophyllea, 144; fimbriata, 143–144; galactina, 141; intybacea, 142; terrestris, 141–143
THIEL, A. F., see STAKMAN, E. C.
Thielavia basicola, 433; breeding tobacco resistant to, 49; on bean seedlings, 63
THOMPSON, NOEL F., The effect of certain chemicals especially copper sulphate and sodium chloride on the germination of bunt spores, 37–38
Thuja plicata, host of Thelephora terrestris, 142; plicata, host of Polyporus schweinitzii, 176; plicata, host of Poria weirii, 176; plicata, host of Fomes pini, 176; plicata, host of Fomes annosus, 176; plicata, host of Polyporus sulphureus, 176; plicata, host of Armillaria mellea, 176; occidentalis, sap stained by Ceratostomella sp., 214–224
THURSTON, H. W. JR., A note on the corrosive sublimate treatment for the control of Rhizoctonia, 150–151
Tilletia, laevis, 37–38; tritici, 37–38, 39; triticina, spore load of seed, relation to smut in crop, 103–104; triticina, effect of treatment for, on germination of wheat, 469–484
Tip burn, 56–57, 182–193
TISDALE, ELIZABETH K., see RUMBOLD, CAROLINE
TISDALE, W. H., Two Sclerotium diseases of rice, 42; and JENKINS, J MITCHELL, Rice straighthead and its MAUED control, 42–43; and WILLIAMSON, MAUDE MILLER, Bacterial leaf spot of lima bean, 52
Tissue break-down of cabbage, 44; of citrus fruits, 44; of lettuce, 44
Titration of potato dextrose agar with lactic acid, 491–494
Tobacco, false mildew of, introduced into United States from Dutch East Indies? 430–432; angular spot of, 48–49; host of Thielavia basicola, 49; mosaic of, 47, 388; mosaic, relation of air temperature to, 446–458; seedling disease caused by Fusarium, 49; wildfire, 48–49; wildfire, relation of air temperature to, 446–458
Tolerance to acids by bacteria, 244–250
Tomato, bacterial spot of, 55; foot rot of, 29; fruit rots, 379–380; host of Rhizopus nigricans, 279; leaf curl of, 380–383; leaf spot, control of, influenced by fertilizers, 433–445; mosaic of, 47; mosaic transferred to Solanum dulcamara, S. nigrum, Physalis longifolia, Nicandra physalodes, and Datura stramonium, 47; overwintering of mosaic of, 47; sunburn of, 379–380
Trametes, sp., cultures of, 154; carnea on peach and plum trees, 105; heteromorpha, on wood of Tsuga canadensis, 326; pini, 45; suaveolens, on Salix sp., 326
Trees, falling foliage of, 337–339
Trifolium, medium, bacterial disease of, 50; pratense, bacterial disease of, 50; pratense, mosaic of, 146–147
Triticum dicoccum, 212, durum, 206, 211, 212; vulgare, 206, 211, 212; reaction of F_2 generation (durum x vulgare) to Puccinia graminis, 205–213
TROST, JOHN F. and HOFFER, G. N. Kernel starchiness as an index of susceptibility to root, stalk and ear rots of corn, 33–34
Tsuga canadensis, Hymenochaete tabacina on wood of, 320; canadensis, Polyporus nidulans on wood of, 323 canadensis, Trametes heteromorpha on wood of, 326; heterophylla, host of Polyporus dryadeus, 99; heterophylla host of Thelephora caryophyllea, 144; heterophylla, Fomes applanatus on wood of, 324; heterophylla, Polystictus versicolor on wood of, 321; mertensiana, host of Polyporus dryadeus, 99
Tylenchulus semipenetrans on citrus roots 90–93
Tylenchus tritici, 41

Umbellularia californica, host of Fomes applanatus, 326

Uredo coleosporioides, 45
Uromyces, alchemillae, 388; betae, 388
Ustilgo tritici, incidence in wheat varieties, 507–510; zeae, 32

VALLEAU, W. D., Resistance as a basis for control of corn root rot, 34; Selection of disease-free seed and seed treatments as possible means of contro. of corn root rot, 35
Valsa, sp., causing poplar canker, 45; leucostoma, 236–243, 293; lupini, on Lupinus arboreus, 391; sordida, 163
VAUGHAN, R. E., Inoculated sulphur for potato scab control, 58; see JONES, FRED REUEL
Velvet bean, host of Sclerotium rolfsii, 378
Venturia inaequalis on apple, 43-44
Vicia faba, host of clover mosaic, 147; mosaic of, 202
Violet root rot, 412-423
Viscum, mucilage of, 99

WALKER, J. C., A Macrosporium rot of onion, 53; The occurrence of dodder on onions, 53; Rust of onion followed by a secondary parasite, 87–90; and JONES, L. R., The relation of soil temperature and other factors to onion smut and infection, 52–53
Walnut, see Juglans nigra
Wart of potato, heredity of immunity to, 306–310; character of growth, 57; resistant varieties, 57; status in Pennsylvania, 58
WEBER, G. F. Studies on corn rust, 31-32
WEIMER, J. L., see HARTER, L. L.
WEISS, FREEMAN, and ORTON, C. R., Second report on the reaction of American potato varieties to the wart disease, 57; and HARVEY, R. B. Catalase, hydrogen-ion concentration and growth in the potato wart disease, 57; see ORTON, C. R.
Western, white pine, see Pinus monticola; yellow pine, see Pinus ponderosa
WESTON, W. H. JR., Significant points in the life history of the Philippine maize mildew, 32; A note relative to the recent appearance of the sugar cane downy mildew in the Philippines, 371–375
Weymouth pine, see Pinus strobus
WEIR, JAMES R., The mucilage of mistletoe berries as an adhesive, 99; Polyporus dryadeus (Pers.) Fr. on conifers in the Northwest, 99; Thelephora terrestris, T. fimbriata, and T. caryophyllea on forest tree seedlings, 141–144; Note on Cenangium abietis (Pers.) Rehm on Pinus ponderosa Laws., 166–170; Polyporus schweinitzii Fr. on Thuja plicata, 176; Cenangium **piniphilum** n. sp., an undescribed canker forming fungus on Pinus ponderosa and P. contorta, 294–296
WELLES, COLIN G., see LEE, H. ATHERTON
Wheat, black point of, 515; bunt of, 37–39; foot rot of, 37; host of Wojnowicia graminis, 505–506; injury by seed treatment of, 103; loose smut of, 507–510; orange leaf rust of, 40; reaction to Puccinia graminis, 205–213; scab of, 35, 115–124; seedling blight of, 35–37; stem rust of, 202; stinking smut of, 269–278; take-all of, 37
WHETZEL, H. H., The Phytopathological Society of France, 100–101; and H. B. HUMPHREY, Frederick Kolpin Ravn, biographical sketch and portrait, 1–5
White pine, see Pinus strobus
Wild cucumber, eradication of, 47
Wildfire of tobacco, relation of air temperature to, 446–458
WILLIAMSON, MAUDE MILLER, see JONES, L. R.; see TISDALE, W. B.
Wilting caused by walnut trees, 346
WINELAND, GRACE, see DICKSON, JAMES G.
WINGARD, S. A., see FROMME, F. D.
Witches'-brooms, of Pinus maritima, 27
WOLF, FREDERICK A., and SHUNK, I. V Tolerance to acids of certain bacterial plant pathogens, 244–250; see SHUNK, I. V.
Wood-rotting fungi, field cultures of, 151–

154; new or little-known hosts for, 319–326
Wojnowicia graminis on wheat, 505–506

Yellows, of cabbage, 54; of celery, 54–55
YORK, H. H., see PENNINGTON, L. H.
YOUNG, H. C. and BENNETT, C. W., Studies in parasitism in the Fusarium group, 56

ZELLER, S. M., Heart-rot of prune and peach in Oregon, 105; A spur blight of pear caused by Botrytis, 105; ZELLER S. M., and OWENS, C. E. European canker on the Pacific Slope, 464–468
Zea, amylacea, 32; dentata, 32; everta, 32; indurata, 32; ramosa, 32; saccharata 32; tunicala, 32; mays indentata, host of Sphacelotheca reiliana, 515; mays saccharata, host of Sphacelotheca reiliana, 515
Zinc-Bordeaux, 185
ZUNDEL, GEORGE L., Preliminary experiments on injury to wheat from seed treatment in Washington, 103; The effects of treatment for bunt on the germination of wheat, 469–484